

## **Clinical Policy: Triamcinolone ER Injection (Zilretta)**

Reference Number: CP.PHAR.371

Effective Date: 03.01.18

Last Review Date: 02.21

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Triamcinolone acetonide extended-release injectable suspension (Zilretta<sup>®</sup>) is an extended-release synthetic corticosteroid.

### **FDA Approved Indication(s)**

Zilretta is indicated as an intra-articular injection for the management of osteoarthritis pain of the knee.

Limitation(s) of use: The efficacy and safety of repeat administration of Zilretta have not been demonstrated.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Zilretta is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Osteoarthritis of the Knee (must meet all):**

1. Diagnosis of osteoarthritis of the knee;
2. Prescribed by or in consultation with a rheumatologist or an orthopedist;
3. Age  $\geq$  18 years;
4. Failure of  $\geq$  4-week trial of one of the following (a or b), unless contraindicated or clinically significant adverse effects are experienced:
  - a. Oral nonsteroidal anti-inflammatory drug (NSAID) at continuous therapeutic dosing (prescription strength);
  - b. Topical NSAID if member is  $\geq$  75 years old or unable to take oral NSAIDs;  
*\*Prior authorization may be required for topical NSAIDs*
5. Trial of at least one other intra-articular glucocorticoid injection for the knee with a documented positive, but inadequate response (e.g., inadequate pain relief, frequent need of rescue medications such as NSAIDs or opioids, need to decrease or inability to increase activity levels, adequate pain relief but with steroid-induced hyperglycemia);  
*\*Prior authorization may be required for intra-articular glucocorticoids*
6. Dose does not exceed 32 mg as a single intra-articular injection into the knee.

**Approval duration: 3 months (one dose per knee)**

**B. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**II. Continued Therapy**

**A. Osteoarthritis of the Knee**

1. Re-authorization is not permitted. Zilretta is not indicated for repeat administration in the same knee. For an untreated knee, members must meet the initial approval criteria.

**Approval duration: Not applicable**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

NSAID: non-steroidal anti-inflammatory drug

TA: triamcinolone acetonide

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<b>Oral NSAIDs</b>		
diclofenac (Voltaren <sup>®</sup> )	50 mg PO BID to TID	150 mg/day
etodolac (Lodine <sup>®</sup> )	400-500 mg PO BID	1200 mg/day
fenoprofen (Nalfon <sup>®</sup> )	400-600 mg PO TID to QID	3200 mg/day
ibuprofen (Motrin <sup>®</sup> )	400-800 mg PO TID to QID	3200 mg/day
indomethacin (Indocin <sup>®</sup> )	25-50 mg PO BID to TID	200 mg/day
indomethacin SR	75 mg PO QD to BID	150 mg/day
ketoprofen	25-75 mg PO TID to QID	300 mg/day

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
<b>Oral NSAIDs</b>		
meloxicam (Mobic <sup>®</sup> )	7.5-15 mg PO QD	15 mg/day
naproxen (Naprosyn <sup>®</sup> )	250-500 mg PO BID	1500 mg/day
naproxen sodium (Anaprox <sup>®</sup> , Anaprox DS <sup>®</sup> )	275-550 mg PO BID	1650 mg/day
oxaprozin (Daypro <sup>®</sup> )	600-1200 mg PO QD	1800 mg/day
piroxicam (Feldene <sup>®</sup> )	10-20 mg PO QD	20 mg/day
salsalate (Disalcid <sup>®</sup> )	1500 mg PO BID or 1000 mg PO TID	3000 mg/day
sulindac	150 mg-200 mg PO BID	400 mg/day
<b>Topical NSAIDs</b>		
diclofenac 1.5% (Pennsaid <sup>®</sup> )	40 drops QID on each painful knee	160 drops/knee/day
Voltaren <sup>®</sup> Gel 1% (diclofenac)	2-4 g applied to affected area QID	32 g/day
<b>Intra-articular Glucocorticoids</b>		
triamcinolone acetonide (Kenalog <sup>®</sup> )	40 mg (1 mL) for large joints	80 mg/treatment
methylprednisolone acetate (Depo-Medrol <sup>®</sup> )	20-80 mg for large joints	80 mg/treatment

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

#### Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): patients with hypersensitivity to triamcinolone acetonide or any component of the product
- Boxed warning(s): none reported

#### Appendix D: General Information

- Zilretta (extended-release triamcinolone acetonide [TA-ER]) is designed to deliver TA over 12 weeks using extended-release microsphere technology. In contrast, Bodick, et al., 2015, reports that, historically, immediate-release intraarticular glucocorticoids, while demonstrating a large initial analgesic effect, wane over one to four weeks.
- In an evaluation of TA-ER vs immediate-release triamcinolone acetonide (TA-IR) synovial and systemic pharmacokinetics, Krause, et al, 2017, reports that TA-ER demonstrated prolonged residency in the joint (through week 12) relative to TA-IR (through week 6), and consequently showed diminished peak plasma steroid levels relative to TA-IR through week 6. Russell, et al, 2017, reports that in patients with knee osteoarthritis and type-2 diabetes mellitus, TA-ER was associated with a significant and clinically relevant reduction in blood glucose elevation relative to TA-IR 72 hours post-injection.
- In the Zilretta pivotal trial, Conaghan, et al, 2018, reported superiority of TA-ER versus placebo to 12 weeks in average daily pain (ADP) scores (primary endpoint) and continuing TA-ER activity out to 24 weeks. While TA-ER did not show superior

outcomes relative to TA-IR over 12 weeks in ADP scores (secondary endpoint), it was superior to TA-IR at week 12 when evaluated using the exploratory endpoints Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)-A/B/C and Knee injury and Osteoarthritis Outcome Score Quality of Life (KOOS QoL) subscales.

- Conaghan also reports that patients treated with TA-ER used significantly less rescue medication than those treated with TA-IR.
- A phase 3b, open-label, single-arm study by Spitzer et al., 2019, evaluated the safety and efficacy of repeat administration of Zilretta in 208 patients, of whom 179 received a second injection of Zilretta after a median of 16.6 weeks. Additional injections after the second dose were not allowed.
  - The proportion of patients who experienced arthralgia in any joint was nearly doubled during the second injection period (19.0%) compared to the first injection period (10.6%); there were also slightly higher rates of index-knee treatment-emergent AEs during the second injection period (17.3%) compared to the first (14.0%).
  - The FDA highlights this concern in the Zilretta Prescribing Information, Section 6.1 Adverse Reactions – Clinical Studies, stating “The data from this study are insufficient to fully characterize the safety of repeat administration of Zilretta.” As a result, the label continues to retain a limitation of use concerning the unknown benefit of repeat administration.

#### V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Osteoarthritis of the knee	32 mg (5 mL) as a single intra-articular extended-release injection	32 mg (5 mL)

#### VI. Product Availability

Injectable suspension of microspheres (single-dose vial for reconstitution): 32 mg/5 mL

#### VII. References

1. Zilretta Prescribing Information. Burlington, MA: Flexion Therapeutics, Inc.; January 2020. Available at: <http://www.zilrettalabel.com/PI.pdf>. Accessed March 26, 2021.
2. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2019. Available at: <https://www.clinicalpharmacology-ip.com>.
3. Hochberg MC, Altman RD, April KT, et al. American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care & Research*. 2012; 64(4): 465-474.
4. Brown GA. American Academy of Orthopaedic Surgeons clinical practice guidelines: Treatment of osteoarthritis of the knee: Evidence-based guideline, 2nd edition. *J Am Acad Orthop Surg*. 2013;21(9):577-9. doi: 10.5435/JAAOS-21-09-577.
5. McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage*. 2014; 22:363-388.
6. Bodick N, Lufkin J, Willwerth C, et al. An intra-articular, extended-release formulation of triamcinolone acetonide prolongs and amplifies analgesic effect in patients with osteoarthritis of the knee: a randomized clinical trial. *J Bone Joint Surg Am*. 2015; 97: 877-88. <http://dx.doi.org/10.2106/JBJS.N.00918>

7. Nelson AE, Allen KD, Golightly YM, et al. A systematic review of recommendations and guidelines for the management of osteoarthritis: The chronic osteoarthritis management initiative of the U.S. Bone and Joint Initiative. *Semin Arthritis Rheum.* 2014; 43:701-712.
8. Rannou F, Peletier JP, Martel-Pelletier J. Efficacy and safety of topical NSAIDs in the management of osteoarthritis: Evidence from real-life setting trials and surveys. *Semin Arthritis Rheum.* 2016; 45:S18-S21.
9. Russell SJ, Sala R, Conaghan PG, et al. In type 2 diabetes mellitus patients with knee osteoarthritis intra-articular injection of FX006 (Extended Release Triamcinolone) is associated with reduced blood glucose elevation vs. standard triamcinolone; a randomized, blinded, parallel group study. *Diabetes.* 2017; 66(Suppl 1): A289.
10. Conaghan PG, Hunter DJ, Cohen SB, et al. Effects of a single intra-articular injection of a microsphere formulation of triamcinolone acetonide on knee osteoarthritis pain. A double-blind, randomized, placebo controlled, multinational study. *J Bone Joint Surg Am.* 2018; 100(8): 666-677.
11. Krause VB, Conaghan PG, Aazami HA, et al. Synovial and systemic pharmacokinetics (PK) of triamcinolone acetonide (TA) following intra-articular (IA) injection of an extended release microsphere-based formulation (FX006) or standard crystalline suspension in patients with knee osteoarthritis (OA). *Osteoarthritis and Cartilage.* 2018; 26: 34-42.
12. Spitzer AI, Richmond JC, Kraus VB, et al. Safety and efficacy of repeat administration of triamcinolone acetonide extended-release in osteoarthritis of the knee: A phase 3b, open-label study. *Rheumatol Ther.* Published online February 11, 2019. <https://doi.org/10.1007/s40744-019-0140-z>.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3304	Injection, triamcinolone acetonide, preservative-free, extended-release, microsphere formulation, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.09.18	02.18
No significant changes; per SDC decision, added HIM Medical Benefit line of business.	04.12.18	
1Q 2019 annual review; no significant changes; references reviewed and updated.	12.11.18	02.19
1Q 2020 annual review: no significant changes; modified NSAID trial duration to 4 weeks to align with existing requirements for hyaluronates; replaced HIM Medical Benefit with HIM line of business; references reviewed and updated.	11.26.19	02.20

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2021 annual review: no significant changes; references to HIM.PHAR.21 revised to HIM.PA.154; added coding implications; references reviewed and updated.	10.22.20	02.21
Added information regarding repeat administration to Appendix D.	03.26.21	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.